

VERIFY™

**LONG-LASTING RESIDUAL ACTIVITY AGAINST GRASS
AND SMALL-SEEDED BROADLEAF WEEDS.**



Verify™ herbicide is for control of yellow nutsedge and the annual grasses and broadleaf weeds listed in the WEEDS CONTROLLED section of the product label. This product alone will not control emerged seedlings. Verify may be applied either as a surface application before or after planting, or after crop emergence. It may also be shallowly incorporated prior to planting to blend the herbicide treatment into the upper 1 to 2 inches of soil. Except for minimum or conservation tillage systems, the seedbed should be fine, firm, and free of clods and trash.

PRODUCT FEATURES

- Provides consistent, season-long control of grass and small-seeded broadleaf weeds, including barnyardgrass, lambsquarters, common waterhemp, henbit, and foxtail species.
- Can be applied preplant, preemergence, and postemergence (up to 11" corn) in single or split applications.
- Excellent tank-mix partner with atrazine, dicamba, simazine, and imazethapyr to broaden the spectrum of control.
- Labeled for use on corn, Miscanthus, and non-food perennial bioenergy crops.

See label for complete application guidelines and restrictions.

ACTIVE INGREDIENT

7 lb/gal acetochlor

HRAC GROUP

15

FORMULATION

Emulsifiable concentrate

REI

12 hours (see exception for soil injection & soil-incorporated uses)

COMPARE TO

Harness®

MODE OF ACTION

Long-chain fatty acid inhibitor

SIGNAL WORD

Warning

RESTRICTED USE

No

PACKAGING

NR 265, bulk

EPA REG NUMBER

83979-8 | AD111623

FREQUENTLY ASKED QUESTIONS

What is the maximum annual use rate for Verify herbicide?

- Maximum annual acetochlor broadcast application rates for all listed crops must not exceed 3.0 pounds active ingredient (3.4 pints of Verify) per acre. Note: One pint per acre of Verify delivers 0.875 pound active ingredient acetochlor per acre.

How should Verify be applied for best results?

Early pre-plant surface

- This product and some labeled tank mixtures of this product may be applied in no-till and other conservation tillage systems before weeds emerge and up to 45 days before planting field corn or silage corn. Split applications can be made 30 to 45 days prior to planting with 60 percent of the specified broadcast rate applied initially and the remaining 40 percent applied at planting. Applications made less than 30 days prior to planting can be made either as a split or a single application.
- Refer to product label for preplant incorporated application directions and use restrictions.

Preemergence surface application

- Verify and all labeled tank mixtures may be applied to the soil surface after planting and prior to either crop or weed emergence. Apply within 5 days of last pre-plant tillage. If weeds emerge after treatment, or if treatment is applied more than 5 days after last pre-plant tillage, rotary hoe or shallowly cultivate immediately to improve performance.
- Precipitation or overhead sprinkler irrigation is required after application to move the herbicide treatment into the weed germination zone. Refer to label for precipitation/irrigation requirements.

Postemergence surface application

- Verify and certain tank mixtures may be applied postemergence until corn reaches 11 inches in height. Application must be made prior to weed seedling emergence or in a tank-mixture that controls emerged weeds. Refer to DIRECTIONS FOR USE on product label for tank mix recommendations, growth stage limitations, and directions.
- Precipitation or overhead sprinkler irrigation is required after application to move the herbicide treatment into the weed germination zone to control unemerged weeds. Refer to label for precipitation/irrigation requirements.

How should Verify be used to control emerged weeds?

- In early preplant applications if emerged weeds are present at the time of treatment, a glyphosate agricultural herbicide, Gramoxone® brand herbicide or 2,4-D should be added to this product according to the directions for use on their respective product labels. Weeds emerged at the time of application are not controlled by this product. If weeds are emerged at application, shallowly cultivate or rotary hoe to improve performance.
- Extreme care must be used when applying tank mixtures to prevent injury to desirable plants and crops



◀ See label for complete application guidelines and restrictions.

©2024 Albaugh, LLC. All rights reserved. Albaugh and the Albaugh logo are registered trademarks and Verify is a trademark of Albaugh, LLC. Harness is a registered trademark of Monsanto Technology LLP. Gramoxone is a registered trademark of a Syngenta Group Company. Always read and follow label restrictions, precautions and directions for use. Some products may not be registered for sale or use in all states or counties. Please check with your local extension service to ensure registration status.