

# BROX<sup>®</sup> 2EC

**FOR POSTEMERGENCE CONTROL OF A VARIETY OF BROADLEAF WEEDS IN CROP AND NON-CROP APPLICATIONS.**

Brox<sup>®</sup> 2EC is a selective postemergence herbicide for control of important broadleaf weeds infesting corn (field and pop), sorghum (grain and forage), wheat, barley, oats, rye, triticale, alfalfa (seedling), flax, onions, garlic, mint (established peppermint and spearmint), Conservation Reserve Program (CRP) areas, grasses grown for seed or sod production, non-residential turfgrass, and non-cropland and industrial sites.

Optimum weed control is obtained when Brox 2EC is applied to actively growing weed seedlings. It is primarily a contact herbicide; therefore, thorough coverage of the weed seedlings is essential for optimum control. Brox 2EC has little residual activity. Therefore, subsequent flushes of weeds will not be controlled by the initial treatment. Generally, crops that form a good canopy will help shade subsequent weed flushes. However, certain crops or short-straw varieties may not develop the crop canopy fast enough to shade the subsequent flushes of weeds.

## PRODUCT FEATURES

- May be used in many crop and non-crop applications providing effective broadleaf weed control.
- Provides excellent tank mix compatibility with a wide array of herbicides, insecticides, fertilizer, and plant growth regulator products.
- Flexible application by ground, aerial or chemigation methods.

See label for complete application guidelines and restrictions.



### ACTIVE INGREDIENT

2.0 lbs/gal bromoxynil octanoate

### HRAC GROUP

6

### FORMULATION

Emulsifiable concentrate

### REI

24 hours (12 hours on sod farm turf)

### COMPARE TO

Buctril

### MODE OF ACTION

Photosynthesis inhibitor

### SIGNAL WORD

Warning

### RESTRICTED USE

No

### PACKAGING

2 x 2.5 gal, 30 gal, NR 265

### EPA REG NUMBER

42750-48 | AD012523

# BROX<sup>®</sup> 2EC

## FREQUENTLY ASKED QUESTIONS

### Can I tank mix Brox 2EC with other crop protection products?

- Yes, Brox 2EC can be applied in tank mixture with other pesticide products registered for use on approved crops provided that these other products are registered for use on the crop/use site to be treated. *Refer to label pages for tank mixing procedures and a list of insecticide, herbicide and plant growth regulator products that are compatible with Brox 2EC.*

### How should I apply Brox 2EC by ground for best results?

- Use a standard herbicide boom sprayer that provides uniform and accurate application. Sprayer should be equipped with screens no finer than 50 mesh in the nozzle tips and in-line strainers.
- For optimum spray distribution and thorough coverage, use of flat-fan nozzles (maximum tip size 8008) with a minimum spray pressure of 30 psi at the nozzle tips are recommended.
- In general, a minimum spray volume of 10 gallons per acre (GPA) is recommended for optimum spray coverage. A minimum of 5 GPA with a minimum spray pressure of 50 psi may be used with higher speed, low-volume ground application if ground terrain, crop, and weed density allow effective spray distribution.
- When using higher speed equipment, a maximum ground speed of 10 mph is suggested if field conditions cause excessive boom movement during application and subsequent poor spray coverage.
- *Refer to label for additional information on ground, aerial and sprinkler irrigation application procedures.*



◀ See label for complete application guidelines and restrictions.

©2023 Albaugh, LLC. All rights reserved. Albaugh, the Albaugh logo and Brox are registered trademarks of Albaugh, LLC. Always read and follow label restrictions, precautions and directions for use. Some products may not be registered for sale or use in all states or counties. Please check with your local extension service to ensure registration status.