

CAPTAN 80WDG

MULTI-SITE CONTACT FUNGICIDE FOR THE CONTROL OF A VARIETY OF DISEASES IN A WATER-DISPERSIBLE GRANULE FORMULATION.



Captan 80 WDG is a multi-site contact fungicide that rapidly penetrates fungal spores and can be used as an effective protectant in a variety of crops including apples, almonds, grapes, pears, cherries, strawberries, and ornamentals. Captan 80 WDG is most effective when used in a protective schedule before infection occurs.

PRODUCT FEATURES

- Excellent multi-site contact fungicide that provides control of diseases such as *Phomopsis*, scab, leaf spot, powdery mildew, brown rot, *Botrytis* (blossom-end rot & bunch rot) and many others.
- Highly concentrated, dry formulation allows for efficient storage and handling.
- Applications may be made by ground or air providing residual control of a variety of diseases.
- May be applied with other EPA approved fungicides for improved disease control and as part of an integrated pest management program.

ACTIVE INGREDIENT 80% captan

FRAC GROUP

FORMULATION Water dispersible granule

REI 24-48 hours, depending on crop

COMPARE TO N/A

MODE OF ACTION Multi-site: phthalimide fungicide

SIGNAL WORD
Danger

RESTRICTED USE

PACKAGING 30 lb bag

EPA REG NUMBER 42750-231 | AD082117



See label for complete application guidelines and restrictions.

CAPTAN 80WDG

FREQUENTLY ASKED QUESTIONS

How should I use Captan 80 WDG on apples for best results?

East of the Rocky Mountains

- Apply 2.5-5.0 lbs./A for control of labeled diseases, following application instructions shown on the product label.
- Do not use in combination with or closely following or in alternation with wettable sulfur products on sulfur sensitive varieties of apples such as Red Delicious, Staymen, Baldwin, King, etc., as severe injury and defoliation may occur.
- Do not apply more than 40 pounds of Captan 80 WDG per acre per year.
- May be applied up to day of harvest.

West of the Rocky Mountains

- Apply 2.5-5.0 lbs./A for control of labeled diseases, following application instructions shown on the product label.
- Do not apply more than 40 pounds of Captan 80 WDG per acre per year.
- May be applied up to day of harvest.

Pacific Northwest

- Apply 3.75 lbs./A for control of labeled diseases, following application instructions shown on the product label.
- Do not apply more than 40 pounds of Captan 80 WDG per acre per year.
- May be applied up to day of harvest.

How should I use Captan 4L on strawberries for best results?

- Apply 1.87-3.75 lbs./A for control of Botrytis (gray mold) and leaf spot, following application instructions shown on the product label.
- Apply by broadcast in sufficient water for thorough coverage by ground or in 10 to 20 gallons of water by air.
- Begin applications when new growth starts in the spring and before fruit starts to form. Repeat at 7- to 14- day intervals.
- Do not apply more than 30 pounds of Captan 80 WDG per acre per year.
- May be applied up to day of harvest.



See label for complete application guidelines and restrictions.

©2023 Albaugh, LLC. All rights reserved. Albaugh and the Albaugh logo are registered trademarks of Albaugh, LLC. Always read and follow label restrictions, precautions and directions for use. Some products may not be registered for sale or use in all states or counties. Please check with your local extension service to ensure registration status.

