

AnchorTM 3LSTTM

TREATED SEED

This seed has been treated with:
Mefenoxam fungicide
(ANCHORTM 3L ST, EPA Reg. 42750-390)

lb AI/100 lb Seed	mg AI per seed	Active Ingredient
_____	OR _____	Mefenoxam

CAUTION

DO NOT USE TREATED SEED FOR FEED, FOOD, OR OIL PURPOSE.

DO NOT allow children, pets, or livestock to have access to treated seed.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Wear long-sleeved shirt, long pants, shoes, socks, and chemical-resistant gloves when handling treated seed.

FIRST AID

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. **DO NOT** induce vomiting unless told to do so by the poison control center or doctor. **DO NOT** give any liquid to the person. **DO NOT** give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center (888-347-6732) or doctor, or going for treatment.

ENVIRONMENTAL HAZARDS

Treated seed exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading and planting, such as in row ends. Dispose of excess treated seed by burying seed away from bodies of water. **DO NOT** contaminate water bodies when disposing of planting equipment washwater.

Groundwater Advisory: Mefenoxam is known to leach through soil into groundwater under certain conditions as a result of label use. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

Surface Water Advisory: This product may impact surface water quality due to runoff of rainwater. This is especially true for poorly draining soils and soils with shallow groundwater. This product is classified as having high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained buffer strip between areas to which this product is applied and surface water features including ponds, streams, and springs will reduce the potential loading of mefenoxam from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

RESTRICTIONS

Treated seed must be incorporated into the soil at the recommended depth.

The maximum application rate for metalaxyl or mefenoxam must not exceed 12.3 lb ai/A per calendar year or the lesser value listed in the crop use section.

Maximum usage when applying both mefenoxam- and metalaxyl-containing products to the same crop within the same season: **DO NOT** apply more than the maximum yearly total application rate for the active ingredient as stated on the label of the product containing the lowest yearly total on that crop.

SEED STORAGE & DISPOSAL

Store away from feeds and foodstuffs.

DO NOT store leftover treated seed where there is a chance of it becoming mixed with untreated seed.

Dispose of seed packaging in accordance with local requirements.

Excess treated seed may be used for ethanol production only if: (1) By-products are not used for livestock feed, and (2) No measurable residues of pesticides remain in ethanol by-products that are used for agronomic practice.

DO NOT use empty seed bags for any other purpose.

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE
Call CHEMTREC (800) 424-9300.

Albaugh, LLC. Ankeny, IA 50021

021723