

TEBUSTAR IM ST

TREATED SEED

This seed has been treated with:

*Metalaxyl fungicide, Tebuconazole fungicide and Imidacloprid insecticide
(EPA Reg. No. 42750-207)*

lb AI/100 lb Seed		mg AI per seed	Active Ingredient
_____		_____	Metalaxyl
_____	OR	_____	Tebuconazole
_____		_____	Imidacloprid

CAUTION

DO NOT USE FOR FEED, FOOD, OR OIL PURPOSE.

DO NOT allow children, pets, or livestock to have access to treated seed.

User is responsible for ensuring that the seed bag meets all requirements under the Federal Seed Act.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Wear long-sleeved shirt, long pants, and chemical-resistant gloves when handling the treated seed.

FIRST AID

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. **DO NOT** induce vomiting unless told to do so by the poison control center or doctor. **DO NOT** give any liquid to the person. **DO NOT** give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center (888-347-6732) or doctor, or going for treatment.

ENVIRONMENTAL HAZARDS

This pesticide is highly toxic to birds and aquatic invertebrates and toxic to fish. **DO NOT** apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark.

Exposed treated seeds may be hazardous to birds. In the event that treated seeds are spilled and cannot be salvaged, completely cover them with soil.

Dispose of all excess treated seed. Leftover treated seed may be double-sown around the headland or buried away from bodies of water in accordance with local requirements.

DO NOT contaminate water bodies when disposing of planting equipment wash waters.

Groundwater Advisory: Metalaxyl is known to leach through soil into groundwater under certain conditions as a result of label use. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

RESTRICTIONS

Treated seed must be planted into the soil at a depth greater than 1 inch.

When applying both metalaxyl-containing and mefenoxam-containing products to the same crop within the same season do not apply more than the maximum yearly total application rate for the active ingredient as stated on the label of the product containing the lowest yearly total on that crop.

Treated areas may be replanted with any crop specified on the TebuStar IM ST label, or for any crop which a tolerance exists for Imidacloprid, as soon as practical following the last application. The following plant-back intervals must be observed for listed crops:

Immediate Plant-Back: Artichoke, Barley, Brassica (cole), Leafy Vegetables, Canola, Cilantro, Corn (field), Corn (sweet), Cotton, Cranberry, Cucurbits, Eggplant, Ground cherry, Leaf petiole vegetable, Legume vegetables (succulent or dried, including soybean), mustard seed, okra, pepinos, pepper, popcorn, potato, rapeseed, sorghum, strawberry, sugar beet, tomatillo, tomato, triticale, turnip greens, vegetable root and tubers, watercress, wheat

30-Day Plant-Back: Cereals (including buckwheat, millet, oats, popcorn, rice, and rye), safflower

Cover crops for soil rebuilding or erosion control may be planted any time, but do not graze or harvest for food or feed.

All other crops not listed on an Imidacloprid label, or for crops for which no tolerance for the active ingredient has been established, a 12-month plant-back interval must be observed.

SEED STORAGE & DISPOSAL

Store treated seed in cool, dry conditions away from feeds and foodstuffs.

DO NOT store leftover treated seed where there is a chance of it becoming mixed with untreated seed.

DO NOT store treated seed in burlap bags or impervious bags/container or in areas that are poorly ventilated.

Dispose of seed packaging in accordance with local requirements.

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE

Call CHEMTREC (800) 424-9300.

Albaugh, LLC. Ankeny, IA 50021

032023