

**STOP! READ THE LABEL AND PAMPHLET BEFORE USING THE PRODUCT, CONSULT
A PROFESSIONAL IN AGRICULTURAL SCIENCES**

SADDLER® 35 FS

**INSECTICIDE – CARBAMATE
THIODICARB**



DANGER

**ANTIDOTE: ATROPINE SULFATE
DENSITY: 1.118 g/ml to 25 °C**

**THIS PRODUCT MAY BE LETHAL IF SWALLOWED AND/OR INHALED, IT MAY CAUSE
DAMAGE TO EYES AND SKIN BY EXPOSURE**

**DO NOT STORE DWELLINGS
MANTAIN AWAY FORM CHILDREN, MENTALLY ILL PEOPLE,
DOMESTIC ANIMALS, FOOD AND MEDICINE**



AGRONOMIC USE: Insecticide, Seed treatment

MODE OF ACTION:

SADDLER® 35 FS is an insecticide for application to the seeds, once the seed is planted at the final site; the active ingredient is slowly liberated controlling the present pests by contact and/or ingestion. Once the seeds germinate, the roots absorb the insecticide, giving protection to the foliage. The product is orient to the seed treatment therefore it is very important to check that good coverage is obtained.

APPLICATION EQUIPMENT:

Place the seed to be treated (clean without dust, straw rest or any other object) in a mixer machine for seeds or in a plastic bag. Add the recommended dosage of **SADDLER® 35 FS** (it can be apply undiluted or it can be mixed with water, the volumen of water has to be the same than the volumen of **SADDLER® 35 FS**). You can also place the seed extended over a plastic on the ground and spray it with a backpack using high volumen nozzles (8004 or 8005). Ensure application equipment is clean and in good condition. **SADDLER® 35 FS** does not have any harmful action on the application equipment. Use the following protective equipment to handle the product during the preparation of the mixture, charging and application: rubber boots, face mask, goggles and chemical-resistant gloves.



PREPARING THE MIXTURE:

If it is considered necessary, you can mix **SADDLER® 35 FS** with a similar amount of water to improve the coverage of the seed. For a better distribution of the product on the seed, let the mixer work or agite the palstic bag for 3 to 5 minutes. Let the seed dry on the shade and then, proceed with sowing. Treat only the seed you will use during the day.



RECOMMENDED USES: USES AUTORIZED IN BELIZE

CROPS	INSECTS CONTROLLED	DOSAGE
Maize (<i>Zea mays</i>)	Fallarmy worm (<i>Spodoptera frugiperda</i>), Leaf beetle (<i>Diabrotica</i> spp.), Army worm (<i>Spodoptera exigua</i>), False meter (<i>Trichoplusia ni</i>), May beetles (<i>Phyllophaga</i> spp.), Wire worm (<i>Agriotes</i> spp.), Cutworm (<i>Agrotis</i> spp.)	1 Liter per 46 Kg of seed
Sorghum (<i>Sorghum bicolor</i>)		
Bean (<i>Phaseolus</i> spp.)		
Trigo (<i>Triticum aestivum</i>)		
Soybean (<i>Glycine max</i>)		
Cotton (<i>Gossypium hirsutum</i>)		
Sesame (<i>Sesamun indicum</i>)		

APPLICATION TIMING:

SADDLER® 35 FS is use only for seed treatment, do not use it foliar applications.

PRE - HARVEST INTERVAL (PHI):

SADDLER® 35 FS is apply just one time as a seed treatment.

RE - ENTRY INTERVAL:

It is recommended to re-enter once the seed is cover by the soil.

PHYTOTOXICITY:

SADDLER® 35 FS is not phytotoxic to crops at the recommended rates given in this pamphlet.

COMPATIBILITY:

SADDLER® 35 FS is compatible and it can be applied in mix with fungicides for seed treatment, but before making a mix for the first time it is recommended to make a small scale test.

PRECAUTIONS AND INSTRUCTIONS OF USE:

STORAGE AND TRANSPORT:

Do not transport or store this product together with food, medicine, clothing, household utensils or forages. Store this product under lock, in a fresh, dry place, away from heat or fire. Keep and store this product only in its original container, with corresponding label and well sealed. Transport and store away from herbicides. **SADDLER® 35 FS** is not flammable or corrosive, but do not place near to sources of high heat.

**DO NOT STORE IN DWELLINGS
MANTAIN AWAY FROM CHILDREN, MENTALLY ILL PERSONS DOMESTIC
ANIMALS, FOOD AND MEDICINE**



**DO NOT EAT, SMOKE OR DRINK DURING THE HANDLING AND
APPLICATION OF THIS PRODUCT, TAKE A SHOWER AND PUT ON
CLEAN CLOTHS AFTER WORK**



SYMPTOMS OF INTOXICATION:

Symptoms of intoxication include midrise (pupil dilatation), ataxia (dizziness) and muscular tremors.

FIRST AID:

BY INGESTION: Do not induce vomiting. Seek medical attention. Give the doctor the label or pamphlet of this product.

BY CONTACT WITH SKIN: If there is contact with skin, rinse with abundant water and soap. Remove all contaminated cloths and wash affected area with water and soap.

BY INHALATION: Move the affected person to a place with fresh air and maintain at rest until completely recovered.

BY CONTACT WITH EYES: Rinse affected eye with abundant fresh water for 15 minutes. If the irritation continues, seek medical attention.

NEVER GIVE TO DRINK NOR INDUCE VOMITING TO A PERSON IN AN UNCONSCIOUS STATE

ANTIDOTE AND MEDICAL TREATMENT:

The antidote is atropine, in all cases of severe intoxication apply preferably by intravenous injection, atropine sulfate at rates of 2-4 mg per adult (0.04 to 0.08 mg/kg of body weight for children) and repeat the dosage of 2-4 mg for adults (0.04 to 0.08 mg/kg for children) each 3-10 minutes until adequate level of atropine is reached, visible by dilatation of the pupils and red color the mouth. If necessary, apply artificial respiration.

NATIONAL CENTERS OF INTOXICATIONS

COUNTRY	INSTITUTE	TELEPHONE
Belize	Centro de Intoxicaciones Karl Heusner Memorial Hospital	(501) 223-1548

NORMS TO PROTECT THE ENVIRONMENT

TOXIC TO FISH AND CRUSTACEAE



DO NOT CONTAMINATE RIVERS, LAKES, PONDS OR GROUND WATER WITH THIS PRODUCT OR WITH CONTAINERS AND BOTTLES

MANAGEMENT OF BOTTLES, PACKING, RESIDUES AND REMANENTS

In case of liquid spillage, leaks or other residues of pesticides, cover and absorb with sand, wood chips or any absorbent material. Put the residues in a hermetic sealed container and take to the distributor or eliminate them in a land fill authorized by the appropriate authority.

Use the entire content of the bottles and when empty wash and triple rinse with clean water, adding the water from the triple rinse back to the spray tank. Destroy or puncture the empty containers and if the country has an official program for collection of empty container disposal, take them to the closest collection center or dispose according with the instructions of the distributor of the product.

EMPTY CONTAINERS NEED TO BE WASHED THREE TIMES; WATER RESULTING FROM THE TRIPLE RINSE MUST BE ADDED TO THE SPRAY TANK.

USING CONTAINERS AND PACKAGES FOR PURPOSES OTHER THAN THOSE, FOR WHICH THEY WERE DESIGNED, MAY BE HAZARDOUS TO HUMAN HEALTH AND THE ENVIRONMENT.

GUARANTEE NOTE

The manufacturer and registrant guarantees the quality of the product in its original and hermetically sealed container. It also guarantees the efficacy of the product, if it is used as recommended in the pamphlet. Since the storage and handling is out of our control, the buyer assumes any risks to persons and property arising from the improper use or handling of this product and accepts the product under these conditions.

FORMULATED BY:

JIANGSU ROTAM CHEMISTRY, CO., LTD. CHINA.
Nº 88 Rotam Road, ETDZ, Kunshan 215301, Jiangsu.
TEL: 0512 – 5771 – 1988 FAX: 0512 – 5771 – 8697

DISTRIBUTOR:

PROSSER FERTILIZER AND AGROTEC CO. LTD
Mile 7.5, George Price Hwy, Belize.

COUNTRY AND REGISTRATION NUMBER

BELIZE 0225-3