

TT-ICASSO 24EC-LEAFLET-20190802

SIZE:280(W)X150(H)MM PTTI12M010190802

PRECAUTIONS AND WARNING FOR USE:

·Do not store in dwellings. Keep under lock and key, away from the reach of children.

·Use the following protective gear when handling or applying this product: rubber boots, respirators, goggles and gloves.

·Do not eat smoke or drink while handling or applying this product.

·Shower after working with this product and wear clean clothes. Do not store and Transport next to food or animal feed.



IN CASE OF INTOXICATION

TAKE THE PATIENT TO A PHYSICIAN AND PRESENT THIS LABEL

SYMPTOMS OF INTOXICATION:

Eyes, nose, throat and skin irritation. If ingested can cause gastroenteritis, nausea, diarrhea, convulsions and heartburn.

FIRST AID:

INGESTION: Do not induce vomiting and immediately transfer patient to the nearest hospital as soon as possible.

CONTACT WITH SKIN: Remove contaminated clothing and wash affected skin with plenty of soap and water.

INHALATION: Seek fresh air immediately. Get medical help if breathing is not normal.

CONTACT WITH EYES: Immediately flush eyes with plenty of water for 15 minutes. Seek medical attention.

DO NOT GIVE TO DRINK OR INDUCE VOMITING TO PERSONS IN AN UNCONCIOUS STATE

MEDICAL TREATMENT: No specific antidote exists. Follow symptomatic therapy.



ICASSO® 24 EC

Clethodim - Herbicide

CHEMICAL COMPOSITION

	% w/w
Active Ingredient	
Clethodim	24%
Inert Ingredients.....	76%
TOTAL.....	100%
Content 240 grams of active ingredient per liter of commercial product	

**READ INSTRUCTIONS BEFORE USE
KEEP OUT OF THE REACH OF CHILDREN**

FORMULATED BY:

JIANGSU ROTAM CHEMISTRY CO. LTD., CHINA
TEL: 0512 – 5771 – 1988 FAX: 0512 – 5771 – 8697

DISTRIBUTED BY:

MAFAS LTD

Lot 167 Concord Rd, BayView, La Romain
Trinidad, W.I. 1-868-697-1375

BATCH NO. See Bottle

EXP. DATE: See Bottle

TRINIDAD & TOBAGO REG. NO TTPR1627

PROTECT THE ENVIRONMENT WITH GOOD AGRICULTURAL PRACTICES, COMPLY WITH THE RECOMMENDATIONS PROVIDED IN THE PAMPHLET

DIRECTIONS FOR USE

CROP	GRASS WEEDS CONTROLLED	RATE:L/ha
Tomato Pepper, Melon, Cucumber, Lettuce, Cabbage, broccoli, Pakchoi, Pumpkin	Digitaria (<i>Digitaria</i> spp.) Pig weed (<i>Echinochloa</i> sp.) Goose grass (<i>Eleusine indica</i>) Bahia grass (<i>Paspalum</i> spp.) Itch grass (<i>Rottboellia cochinchinensis</i>) Foxtail (<i>Paspalum maximum</i>)	0.3-0.5 l/ha (2.5-5.0 ml/4.57 L) for annual grasses in broadleaf crops. 0.5-1 L/ha(5-10ml/4.57L) for perennial grasses Mix with sufficient surfactant for better results Apply in early post emergence on actively growing weeds (2-4 leaves).
Melongene, Chive, Celery, Parsley Onion, Field Corn, Bean, Pea, Root Crops, Ornamentals, etc	Jaragua grass (<i>Hypharrenia rufa</i>) Johnsongrass (<i>Sorghum halepense</i>)	

Re-Entry Period 12 hours after application

Method of Application- apply **ICASSO 24 EC** using high or Low volume sprayers. Prepare a pre-mix of the dosage to be used and water, fill spray tank to ½ the volume of required water, stir, add the premix to tank and while stirring top up the rest of the water.

COMPATIBLE with a wide range of agrochemicals

TOXIC TO FISH AND CRUSTACEAE



DO NOT CONTAMINATE RIVERS, LAKES OR PONDS WITH THIS PRODUCT OR ITS EMPTY CONTAINERS OR PACKAGES.

DO NOT RE-USE THIS PACKAGE. Triple rinse empty bottle, apply washings to treatment area, puncture bottle and bury in an approved disposal area

WARRANTY DECLARATION

The seller guarantees the product in this container complies with the terms on the label and pamphlet. The buyer assumes any risks to persons and property arising out of the improper use, storage and handling **ICASSO 24 EC**



WARNING

