

TT-ROTAPRID 35SC-LEAFLET-20190802

SIZE:280(W)X150(H)MM PTTR20M010190802

PRECAUTIONS AND WARNING FOR USE:

Do not store in dwellings. Keep under lock and key, away from the reach of children.

Use the following protective gear when handling or applying this product: rubber boots, respirators, goggles and gloves.

Do not eat smoke or drink while handling or applying this product.

Shower after working with this product and wear clean clothes. Do not store and Transport next to food or animal feed.



IN CASE OF INTOXICATION TAKE THE PATIENT TO A PHYSICIAN AND PRESENT THIS LABEL.

SYMPTOMS OF INTOXICATION:

Eyes, nose, throat and skin irritation. If ingested can cause gastroenteritis, nausea, diarrhea, convulsions and heartburn.

FIRST AID:

INGESTION: Induce vomiting immediately and transfer patient to the nearest hospital as soon as possible.

CONTACT WITH SKIN: Remove contaminated clothing and wash affected skin with plenty of soap and water.

INHALATION: Seek fresh air immediately.

CONTACT WITH EYES: Immediately flush eyes with plenty of water for 15 minutes. Seek medical attention.

DO NOT GIVE TO DRINK OR INDUCE VOMITING TO PERSONS IN AN UNCONSCIOUS STATE.

MEDICAL TREATMENT.

Symptomatic, supply enough fluids if the patient has ingested the product.



ROTAPRID® 35 SC

INSECTICIDE - IMIDACLOPRID

CHEMICAL COMPOSITION:

W/V

Active Ingredients:

1-(6-chloro-3-pyridylmethyl)-N-nitroimidazolidin-2-ylideneamine.....35%
Inert Ingredients65%
TOTAL.....100%

CONTENT: 350 grams of active ingredient per litre of commercial product

**READ INSTRUCTIONS BEFORE USE
KEEP OUT OF THE REACH OF
CHILDREN**

FORMULATED BY:

JIANGSU ROTAM CHEMISTRY CO. LTD., CHINA
TEL: 0512 5771 - 1988 - FAX: 0512 5771 - 8697

DISTRIBUTED BY:

MAFAS LTD
Lot 167 Concord Rd, BayView, La Romain
Trinidad, W.I. 1-868-697-1375

PROTECT THE ENVIRONMENT WITH GOOD AGRICULTURAL PRACTICES, COMPLY WITH THE RECOMMENDATIONS PROVIDED IN THE PAMPHLET

DIRECTIONS FOR USE

CROPS	PESTS	RATE	PHI
Potato	Thrip	0.5-1.0 L Ha 5-10ml/gall	21
Melon	Whitefly	0.5-1.0 L Ha 5-10ml/gall	21
Onion	Aphid	0.5-1.0 L Ha 5-10ml/gall	21
Rice	Leafminer Stinkbugs	0.5-1.0 L Ha 5-10ml/gall	21
Tomato	Whitefly	0.5-1.0 L Ha 5-10ml/gall	21
Citrus	Leaf Hopper	0.5-1.0 L Ha 5-10ml/gall	21

Re-Entry- Once spray has dried completely

Method of Application-Using tractor or backpack sprayers apply at signs of first insect attack, or preventive in fields with history of early attack. Apply to base of the plant from 0-7 days after transplant and repeat 21 days.

COMPATIBLE with a wide range of Agrochemicals

TOXIC TO FISH



DO NOT CONTAMINATE RIVERS, LAKES OR PONDS WITH THIS PRODUCT OR ITS EMPTY CONTAINERS OR PACKAGES.

DO NOT RE-USE THIS PACKAGE

WARRANTY DECLARATION

The seller guarantees the product in this container complies with the terms on the label and pamphlet. The buyer assumes any risks to persons and property arising out of the improper use, storage and handling **ROTAPRID@35 SC**

BATCH NO. See bottle

EXP. DATE: See bottle

TRINIDAD & TOBAGO REG NO. PR 1495

MODERATELY DANGEROUS