

CAPTAN 4L

MULTI-SITE CONTACT FUNGICIDE FOR USE AS AN EFFECTIVE PROTECTANT IN A VARIETY OF LABELED CROPS.



Captan 4L is a multi-site contact fungicide that rapidly penetrates fungal spores and can be used as an effective protectant in a variety of crops including apples, almonds, grapes, cherries, stone fruit, strawberries, and ornamentals. Captan 4L is most effective when used in a protective schedule before infection occurs.

PRODUCT FEATURES

- Excellent multi-site contact fungicide that provides control of diseases such as *Phomopsis*, scab, leaf spot, powdery mildew, brown rot, scab, *Botrytis* (blossom-end rot & Bunch Rot) and many others.
- Applications may be made by ground or air providing residual control of a variety of diseases.
- May be applied with other EPA approved fungicides for improved disease control and as part of an integrated pest management program.

ACTIVE INGREDIENT

3.9 lbs/gal captan

FRAC GROUP

M4

FORMULATION

Aqueous suspension

REI

24-48 hours, depending on crop

COMPARE TO

N/A

MODE OF ACTION

Multi-site: phthalimide fungicide

SIGNAL WORD

Danger

RESTRICTED USE

No

PACKAGING

2 x 2.5 gal

EPA REG NUMBER

42750-236 | AD122017

See label for complete application guidelines and restrictions.

CAPTAN 4L

FREQUENTLY ASKED QUESTIONS

How should I use Captan 4L on apples for best results?

East of the Rocky Mountains

- Apply 3.0-4.0 qt/A for control of primary scab, black rot (frog-eye) and *Botrytis* blossom end rot, following application instructions shown on the product label. Apply at higher rates and shorter intervals when disease pressure is high.
- Do not use in combination with or closely following or in alternation with wettable sulfur products on sulfur sensitive varieties of apples such as Red Delicious, Staymen, Baldwin, King, etc., as severe injury and defoliation may occur. For pre-blossom sprays, use a dilute spray.
- Do not apply more than 32 quarts of Captan 4L per acre per year. *SEE USE PRECAUTIONS ON LABEL FOR DEFINITION OF CROP CYCLE.*
- Preharvest interval (PHI) - May be applied up to day of harvest.

West of the Rocky Mountains

- Apply 3.0-4.0 qt/A for control of apple scab following application instructions shown on the product label. Apply at higher rates and shorter intervals when disease pressure is high. Most effective when used in a protective schedule before infection occurs.
- Do not apply more than 32 quarts of Captan 4L per acre per year.
- May be applied up to day of harvest.

Pacific Northwest

- Apply 4.0 qt/A for control of labeled diseases, following application instructions shown on the product label. Apply at higher rates and shorter intervals when disease pressure is high. Apply 1 or 2 applications with late cover sprays.

- Do not apply more than 32 quarts of Captan 4L per acre per year.
- May be applied up to day of harvest.

How should I use Captan 4L on strawberries for best results?

California

- Apply 1.5-2.5 qt/A for control of *Botrytis* and leaf spot. Apply at higher rates and shorter intervals when disease pressure is high.
- Apply by broadcast in sufficient water for thorough coverage by ground or in 10 to 20 gallons of water by air.
- Begin applications when new growth starts in the spring and before fruit starts to form. Repeat at 7- to 14- day intervals.
- Do not apply more than 24 quarts of Captan 4L per acre per year.
- May be applied up to day of harvest.

U.S., except California

- Apply 1.5-3.0 qt/A for control of *Botrytis* and leaf spot. Apply higher rates and at shorter spray intervals when disease pressure is higher.
- Apply by broadcast in sufficient water for thorough coverage by ground or in 10 to 20 gallons of water by air.
- Begin applications when new growth starts in the spring and before fruit starts to form. Repeat at 7- to 14- day intervals.
- Do not apply more than 24 quarts of Captan 4L per acre per year.
- May be applied up to day of harvest.



◀ See label for complete application guidelines and restrictions.

©2023 Albaugh, LLC. All rights reserved. Albaugh and the Albaugh logo are registered trademarks of Albaugh, LLC. Always read and follow label restrictions, precautions and directions for use. Some products may not be registered for sale or use in all states or counties. Please check with your local extension service to ensure registration status.