

TEBUSTAR[®] 250 ST TREATED SEED

*This seed has been treated with:
Tebuconazole fungicide
(TebuStar 250 ST, EPA Reg. 42750-130)*

lb AI/100 lb Seed	mg AI per seed	Active Ingredient
_____	_____	Tebuconazole
OR		
_____	_____	

CAUTION

DO NOT USE TREATED SEED FOR FEED, FOOD, OR OIL PURPOSE.

DO NOT allow children, pets, or livestock to have access to treated seed.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Wear long pants, long-sleeved shirt, and protective gloves when handling treated seed.

FIRST AID

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. **DO NOT** induce vomiting unless told to do so by the poison control center or doctor. **DO NOT** give any liquid to the person. **DO NOT** give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center (888-347-6732) or doctor, or going for treatment.

ENVIRONMENTAL HAZARDS

Treated seed exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading and planting, such as in row ends. Dispose of all excess treated seed by burying seed away from bodies of water. **DO NOT** contaminate water bodies when disposing of planting equipment washwater.

Groundwater Advisory: Tebuconazole is known to leach through soil into groundwater under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground-water contamination.

Surface Water Advisory: This product may contaminate water through drift of spray in wind. This product has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to runoff that contains this product. A level, well maintained

vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall-runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted within 48 hours.

RESTRICTIONS

Barley, wheat, triticale and oat green forage may be grazed or harvested for hay 31 days after seeding.

Treated seed must be incorporated into the soil at the recommended depth.

SEED STORAGE & DISPOSAL

Store treated seed away from food and feedstuffs.

DO NOT store leftover treated seed where there is a chance of it becoming mixed with untreated seed.

Dispose of seed packaging or containers in accordance with local requirements.

Excess treated seed may be used for ethanol production only if: (1) By-products are not used for livestock feed, and (2) No measurable residues of pesticides remain in ethanol by-products that are used for agronomic practice.

DO NOT use empty seed bags for any other purpose.

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE

Call CHEMTREC (800) 424-9300.

Albaugh, LLC. Ankeny, IA 50021

031723