

THIABENDAZOLE 4L ST TREATED SEED

*This seed has been treated with:
Thiabendazole fungicide (EPA Reg. No. 42750-226)*

lb AI/100 lb Seed	mg AI per seed	Active Ingredient
_____	OR _____	Thiabendazole

CAUTION

DO NOT USE FOR FEED, FOOD, OR OIL PURPOSE.
DO NOT allow children, pets, or livestock to have access to treated seed.

User is responsible for ensuring that the seed bag meets all requirements under the Federal Seed Act.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Wear long-sleeved shirt, long pants, and chemical-resistant gloves when handling the treated seed.

FIRST AID

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. **DO NOT** induce vomiting unless told to do so by the poison control center or doctor. **DO NOT** give any liquid to the person. **DO NOT** give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center (888-347-6732) or doctor, or going for treatment.

ENVIRONMENTAL HAZARDS

Treated seeds exposed on soil surface may be hazardous to wildlife.

Cover or collect treated seeds spilled during loading and planting.

Dispose of all excess treated seed. Leftover treated seed may be double-sown around the headland or buried away from water sources in accordance with local requirements.

This pesticide is toxic to fish. **DO NOT** apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high-water mark. Runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

DO NOT contaminate water bodies when disposing of planting equipment wash water or rinsate.

RESTRICTIONS

Treated seed must be planted into the soil at a depth greater than 1 inch.

DO NOT feed vines or hay grown from treated seed to livestock.

Use crop specific seeding rates according to local agricultural practice.

DO NOT plant any crop, other than soybeans, dry peas, barley, wheat, oats, rye, and triticale, brassica head and stem subgroup 5A, cucurbit vegetable cropgroup 9, root vegetables (except sugarbeets) subgroup 1B, bulb vegetables subgroup 3-07A, spinach and alfalfa, within 30 days to fields in which treated seeds were planted.

Regardless of type of application, **DO NOT** apply more than 0.15 lb thiabendazole per acre (68.0 grams ai/A) per year.

SEED STORAGE & DISPOSAL

Store treated seed away from food and feedstuffs.

DO NOT store leftover treated seed where there is a chance of it becoming mixed with untreated seed.

DO NOT store treated seed in burlap bags or impervious bags/container or in areas that are poorly ventilated.

Dispose of seed packaging or containers in accordance with local requirements.

DO NOT use empty seed bags for any other purpose.

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE
Call CHEMTREC (800) 424-9300.

Albaugh, LLC. Ankeny, IA 50021

032223