

HERBICIDE

TRIAD SFZ SELECT™

SELECTIVE, POSTEMERGENCE COMBINATION HERBICIDE.



Triad SFZ Select™ contains four active ingredients, including sulfentrazone, to broaden the spectrum of postemergence broadleaf weed control with proven performance in turfgrass. Weed injury symptoms can be noticed within hours of the application and plant death can occur within ten to fourteen days. These combined herbicides provide limited residual activity at specified use rates. Sulfentrazone is in the aryl triazolinone family and inhibits protoporphyrinogen oxidase (Protox), a pivotal enzyme in chlorophyll production. Without this key enzyme, a build-up of peroxide-like compounds occur thus causing the plant cell membranes of weeds to rupture.

PRODUCT FEATURES

- Rainfast in 6 hours.
- Visible results in 24-48 hours.
- Water-based formulation.
- Tough on weeds, gentle on established turfgrass.

Tough weed control in industrial or low-maintenance areas

Triad SFZ Select can be used for certain tough-to-control weeds. Mixed stands of Kentucky bluegrass, tall fescue, smooth bromegrass, orchardgrass, and reed canarygrass may be treated. Applications to non-cropland areas (roadsides and rights-of-way) are not applicable to treatment of commercial timber, or other plants being grown for sale, or other commercial use, or for commercial seed production, or for research purposes.

Treatments of **Triad SFZ Select** may injure or kill legumes, including clovers (sweet, yellow, red, crimson, alsike, hop, white), lespedezas, trefoils, and vetches.

LABELED USES

- Residential lawns
 - Golf course turf including fairways, aprons, roughs
 - Sports turf
 - Public recreation areas
- See label for complete list*

See label for complete application guidelines and restrictions.

ACTIVE INGREDIENT

18.33% 2,4-D
6.65% MCPA
2.88% dicamba
0.65% sulfentrazone

HRAC GROUP

2,4-D - Group 4
MCPA - Group 4
Dicamba - Group 4
Sulfentrazone - Group 14

FORMULATION

Liquid

CHEMICAL FAMILY

PPO inhibitor

COMPARE TO

Surge®

MODE OF ACTION

Cell membrane disrupter, causes accelerated growth until death.

RESTRICTED USE

No

SIGNAL WORD

Warning

PACKAGING

2 x 2.5 gal, 5 x 30 gal

EPA REG NUMBER

89442-39

TRIAD SFZ SELECT™

SELECTIVE, POSTEMERGENCE COMBINATION HERBICIDE.

APPLICATION INFORMATION

Spot treatment

With hand operated sprayers (including backpack sprayers, compression sprayers and knapsack sprayers).

- Apply any time the emerged broadleaf weeds are actively growing.
- Calibration and proper application are essential when using this product.
- Uniform applications are essential when using this product. Over-application or rates above those specified on this label, including excessive overlaps of this product can cause turf injury.
- Hand-held techniques. Wands fitted with flat fan nozzle tips may be used with the appropriate technique.
- Follow-up applications as spot treatments at a 30-day interval are suggested for more mature weeds, for dense infestations, and for adverse environmental conditions.
- For cool-season turfgrass: Mix 1.2 to 1.5 fl oz of this product per one (1.0) gallon of water for treatment of approximately 1,000 sq ft of turfgrass. Apply any time the emerged broadleaf weeds are susceptible.
- For warm-season turfgrass: Mix 1.0 to 1.2 fl oz of this product per one (1.0) gallon of water for treatment of approximately 1,000 sq ft of turfgrass. Apply any time the emerged broadleaf weeds are susceptible.
- For centipedegrass, use lower rates within specified range until turfgrass tolerance to injury can be determined.

Limitations on spot treatments for ornamental turfgrass

Spot treatment is defined as a treatment area no greater than 1,000 sq ft per acre. The maximum single application rate is 1.5 fl oz per 1,000 sq ft per application (0.25 lb. MCPA-p acid equivalent per acre). The maximum number of spot treatments is limited to two per year with a minimum of thirty days between applications.

LABELED WEEDS

Aster, white heath and white prairie	Dandelion	Ground ivy	Oxalis (yellow woodsorrel and creeping woodsorrel)	Redweed
Bedstraw	Dayflower	Groundsel	Parsley piert	Shepherd's purse
Beggarweed, creeping	Deadnettle	Hawkweed	Pennsylvania smartweed	Spurge
Black medic	Dock	Heal-all	Pepperweed	Thistle
Broadleaf plantain	Dogfennel	Henbit	Pigweed	Virginia buttonweed
Buckhorn plantain	Dollarweed (pennywort)	Innocence (Blue-eyed Mary)	Pineappleweed	White clover (Dutch clover, honeysuckle clover, white trefoil and purplewort)
Bull thistle	False dandelion (spotted cats ear and common cats ear)	Knotweed	Plantain	Wild carrot
Burdock, common	Field bindweed (morning glory and creeping jenny)	Lambsquarters	Poison ivy	Wild garlic
Buttercup, creeping	Field oxeye daisy (creeping oxeye)	Lawn burweed	Poison oak	Wild geranium
Carpetweed	Filaree, whitestem and redstem	Lespedeza, common	Prickly lettuce (compass plant)	Wild lettuce
Chickweed, common	Florida betony	Mallow, common	Puncturevine	Wild mustard
Chicory	Florida pusley	Matchweed	Purple cudweed	Wild onion
Cinquefoil		Mouseear chickweed	Purslane	Wild strawberry
Clover		Nutsedge (yellow)	Ragweed	Yarrow
Curly Dock		Old world diamond flower	Red sorrel (sheep sorrel)	Yellow rocket



◀ See label for complete application guidelines and restrictions.

©2023 Albaugh, LLC. Albaugh and the A logo are registered trademarks and Triad SFZ Select is a trademark of Albaugh, LLC. Surge is a registered trademark of PBI-Gordon Corporation. Always read and follow label restrictions, precautions, and directions for use. This product may not be registered for sale or use in all states or counties. Please check with your state agency responsible for pesticide registration to ensure registration status.