TRIAD TZ SELECT

BROAD-SPECTRUM WEED CONTROL.



Triad TZ Select[™] contains four active ingredients, with two modes of action to provide dependable postemergence control of broadleaf weeds in a variety of turf types.

Triad TZ Select is quickly taken up by the weed leaves and stems, and weed growth ceases within hours. Weed control effects may be seen within a few hours to a few days after application and death occurs in 1 to 4 weeks after application, depending on the application rate, sensitivity of the weed, and environmental conditions.

PRODUCT FEATURES

- Excellent postemergence activity with proven rapid and effective weed control performance for some of the toughest broadleaf weeds in turfgrass such as dandelion, spurge, white clover, plantains, ground ivy, Virginia buttonweed, and wild violet.
- Labeled for the suppression of yellow nutsedge.
- This product exhibits improved cool-weather performance.
- Can be applied to fully dormant bermuda, zoysia, and bahia grasses.
- The addition of sulfentrazone speeds herbicidal activity.
- This product is generally rainfast in as little as three hours.

LABELED USES

- Residential lawns
- Athletic fields
- Golf courses
- Parks
- Industrial and institutional buildings and more

ACTIVE INGREDIENT

31.82% 2,4-D 8.40% triclopyr 2.43% dicamba acid 0.73% sulfentrazone

HRAC GROUP

2,4-D - Group 4 Triclopyr - Group 4 Dicamba acid - Group 4 Sulfentrazone - Group 14

FORMULATION

Emulsifiable concentrate

CHEMICAL FAMILY

PPO inhibitor (Group 14), Growth regulator (Group 4)

COMPARE TO

 $T\text{-}Zone^{\scriptscriptstyle\mathsf{TM}}$

MODE OF ACTION

Cell membrane disrupters cause accelerated growth until death.

RESTRICTED USE

No

SIGNAL WORD

Caution

PACKAGING

 12×1 qt , 4×1 gal, 2×2.5 gal, 5×30 gal drum

EPA REG NUMBER

89442-52

See label for complete application guidelines and restrictions.



TRIAD TZ SELECT

BROAD-SPECTRUM WEED CONTROL.

APPLICATION INFORMATION

Generally, the lower application rates within the specified range will provide satisfactory control of sensitive weed species. The higher application rates within the specified range will be required for dense infestations of perennial weeds, for adverse/extreme environmental conditions, or for weeds hardened off or more mature.

TABLE 1. Use rates for ornamental turfgrass, sod farms and non-cropland.					
Species	Rate	Spray Volume			
Cool-season turf					
Kentucky bluegrass, annual bluegrass, annual ryegrass, perennial ryegrass, tall fescue, red or fine leaf fescues	3.25 to 4 pints/A (1.2 to 1.5 fl oz/1,000 sq ft)	10 to 220 gal/A (0.25 to 5.0 gal/1,000 sq ft)			
	Warm-season turf (Dormant turf)				
Hybrid Bermudagrass, common Bermudagrass, Zoysiagrass and Bahiagrass	2 to 2.25 pints/A (0.75 to 0.83 fl oz/1,000 sq ft)	10 to 220 gal/A (0.25 to 5.0 gal/1,000 sq ft)			

Dormant turf: This product may be applied to fully dormant bermudagrass, fully dormant zoysiagrass, and fully dormant bahiagrass. **Note: DO NOT** apply to above listed warm-season turfgrass unless turf injury can be tolerated. It is impossible to test all environmental conditions for the listed warm-season turfgrass. We suggest testing this product on a small area and observe the treated area for 30 days to determine the acceptability of turf discoloration.

DO NOT apply this product to warm-season turfgrass during spring green-up or in the fall during the transition period between active growth and dormancy.

TABLE 2. Limitations on broadcast applications to ornamental turfgrass, sod farms and non-cropland.					
Use Site	Maximum rate per application	Maximum number of applications per year*	Minimum interval between applications	Maximum seasonal rate	
Ornamental turfgrass	4.0 pints/A (0.88 lb 2,4-D ae/A) (0.25 lb triclopyr ae/A) (0.10 lb dicamba ae/A) (0.03 lb sulfentrazone ae/A)	2	N/A	8.0 pints/A (1.75 lb 2,4-D ae/A) (0.50 lb triclopyr ae/A) (0.20 lb dicamba ae/A) (0.06 lb sulfentrazone ae/A	
Sod Farms	4.0 pints/A (0.88 lb 2,4-D ae/A) (0.25 lb. triclopyr ae/A) (0.10 lb dicamba ae/A) (0.03 lb sulfentrazone ae/A)	2	21 days	8.0 pints/A (1.75 lb 2,4-D ae/A) (0.50 lb triclopyr ae/A) (0.20 lb dicamba ae/A) (0.06 lb sulfentrazone ae/A	
4.0 pints/A (0.88 lb 2,4-D ae/A) (0.25 lb triclopyr ae/A) (0.10 lb dicamba ae/A) (0.03 lb sulfentrazone ae/A)	1 (woody plants)	N/A	8.0 pints/A (1.75 lb 2,4-D ae/A)		
	(0.25 lb triclopyr ae/A) (0.10 lb dicamba ae/A)	2 (annual and perennial weeds)	30 days	(0.50 lb triclopyr ae/A) (0.20 lb dicamba ae/A) (0.06 lb sulfentrazone ae/A)	

^{*} New York: Only one application per year of this product is allowed.



[■] See label for complete application guidelines and restrictions.

©2023 Albaugh, LLC. Albaugh and the A logo are registered trademarks and Triad TZ Select is a trademark of Albaugh, LLC. T-Zone is a trademark of PBI-Gordon Corporation.

Always read and follow label restrictions, precautions, and directions for use. This product may not be registered for sale or use in all states or counties. Please check with your state agency responsible for pesticide registration to ensure registration status.

